Geometry Solutions

1. Let PQRS be a square with side length 12. Point A lies on segment \overline{QR} with $\angle QPA = 30^{\circ}$, and point B lies on segment \overline{PQ} with $\angle SRB = 60^{\circ}$. What is AB?

Proposed by Gunmay Handa

Answer: $4\sqrt{6}$

Note that since PQ = 12, $PA = 8\sqrt{3}$ so $QA = 4\sqrt{3}$. By similar reasoning and using that $\angle BRQ = 90 - 60 = 30^{\circ}$, $BQ = 4\sqrt{3}$. Then $AB = \sqrt{BQ^2 + QA^2} = \boxed{4\sqrt{6}}$.

2. Let ABC be a triangle. Points D and E are placed on \overline{AC} in the order A, D, E, and C, and point F lies on \overline{AB} with $EF \parallel BC$. Line segments \overline{BD} and \overline{EF} meet at X. If AD=1, DE=3, EC=5, and EF=4, compute FX.

Proposed by Misha Ivkov and David Altizio

Answer: $\frac{5}{8}$

Solution. Since $\triangle AEF \sim \triangle ACB$, $BC = EF \cdot \frac{AC}{AE} = 4 \cdot \frac{9}{4} = 9$. Furthermore, since $\triangle DEX \sim \triangle DCB$, $XE = BC \cdot \frac{DE}{DC} = 9 \cdot \frac{3}{8} = \frac{27}{8}$. Thus $XF = EF - XE = \boxed{\frac{5}{8}}$.

3. Point A, B, C, and D form a rectangle in that order. Point X lies on CD, and segments \overline{BX} and \overline{AC} intersect at P. If the area of triangle BCP is 3 and the area of triangle PXC is 2, what is the area of the entire rectangle?

Proposed by Josh Abrams

Answer: 15

Solution. Let Y be the intersection point of BX with AD. Compute BP: PX = 3:2, and furthermore

$$\frac{3}{2} = \frac{BP}{PX} = \frac{BA}{CX} = \frac{CD}{CX}.$$

So the area of $\triangle XDY$ is $\frac{5}{4}$, and hence the area of $\triangle ABY$ is $3 \cdot \frac{5}{4} = \frac{15}{4}$. Thus the area of ABXD is 10, and we may conclude the answer of $\boxed{15}$.

4. Triangle ABC has a right angle at B. The perpendicular bisector of \overline{AC} meets segment \overline{BC} at D, while the perpendicular bisector of segment \overline{AD} meets \overline{AB} at E. Suppose CE bisects acute $\angle ACB$. What is the measure of angle ACB?

Proposed by Daniel Li

Answer: 36°

Solution. The key observation is that quadrilateral AEDC is cyclic. One can prove this by recalling that E is the midpoint of the arc AD of $\odot(ADC)$. (Alternatively, using the Angle Bisector Theorem yields


$$\frac{BE}{ED} = \frac{BE}{EA} = \frac{BC}{CA},$$

which gives $\triangle DEB \sim \triangle ACB$.)

Now let $\angle ACE = \angle BCE = \alpha$. Then $\angle EAD = \angle ECD = \alpha$ while $\angle DAC = \angle ACD = 2\alpha$; in turn,

$$90^{\circ} = \angle BAC + \angle ACB = 5\alpha.$$

Solving yields $\alpha = 18^{\circ}$, so $\angle ACB = \boxed{36^{\circ}}$



5. For every positive integer k, let $\mathbf{T}_k = (k(k+1), 0)$, and define \mathcal{H}_k as the homothety centered at \mathbf{T}_k with ratio $\frac{1}{2}$ if k is odd and $\frac{2}{3}$ is k is even. Suppose P = (x, y) is a point such that

$$(\mathcal{H}_4 \circ \mathcal{H}_3 \circ \mathcal{H}_2 \circ \mathcal{H}_1)(P) = (20, 20).$$

What is x + y?

Proposed by Gunmay Handa

Answer: 256

Solution. The key claim is that the composition of two homotheties centered at A and B with ratios r and s is a homothety itself with ratio rs and center X lying on AB satisfying $\frac{XA}{XB} = \frac{(s-1)}{s(1-r)}$, where the ratio is directed.

Let P be a point in the plane, P' be the image of P under the homothety at A and P'' be the image of P' under the homothety at B; let $X = P'P'' \cap AB$. Then by Menelaus's theorem on $\triangle P'AB$ and transversal XP, we obtain

$$\frac{PP'}{PA} \cdot \frac{XA}{XB} \cdot \frac{BP''}{P''P} = -1 \implies (1-r) \cdot \frac{XA}{XB} \cdot \frac{s}{1-s} = -1 \implies \frac{XA}{XB} = \frac{(s-1)}{s(1-r)}$$

so the position of X does not depend on P. Moreover, we can apply Menelaus's theorem again on $\triangle PP'P''$ with transversal AB to conclude

$$\frac{XP''}{XP} \cdot \frac{AP}{AP'} \cdot \frac{BP'}{BP''} = 1 \implies \frac{XP''}{XP} = rs$$

so we can conclude the desired.

Using the claim, we conclude the composition of the four homotheties has ratio $\frac{1}{2} \cdot \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{2}{3} = \frac{1}{9}$, and we can compute the center of $\mathcal{H}_2 \circ \mathcal{H}_1$ as (4,0), the center of $\mathcal{H}_4 \circ \mathcal{H}_3$ as (16,0), and the center of the composition of the two aforementioned composited homotheties as (13,0). Hence, x and y satisfy

$$x - 13 = 9(20 - 13)$$
$$y - 0 = 9(20 - 0)$$

so (x, y) = (76, 180) and the answer is 256

6. Two circles ω_A and ω_B have centers at points A and B respectively and intersect at points P and Q in such a way that A, B, P, and Q all lie on a common circle ω . The tangent to ω at P intersects ω_A and ω_B again at points X and Y respectively. Suppose AB = 17 and XY = 20. Compute the sum of the radii of ω_A and ω_B .

Proposed by David Altizio

Answer: $3\sqrt{51}$

Solution. First remark that since PAQB is a kite, $\angle APB = \angle AQB$; combining this with APBQ cyclic implies $\angle APB = \angle AQB = 90^{\circ}$.

The key observation is that P is the midpoint of \overline{XY} . To prove this, let M and N be the feet of the perpendiculars from A and B respectively to XY. Then $\triangle MPA \sim \triangle PBA \sim \triangle NBP$, so

$$PM = PA \cdot \frac{PB}{AB} = PB \cdot \frac{PA}{AB} = PN.$$

Thus P is the midpoint of \overline{MN} , meaning it is also the midpoint of \overline{XY} .

To finish, let r_A and r_B be the radii of the circles ω_A and ω_B respectively. Then $r_A^2 + r_B^2 = 289$ by the Pythagorean Theorem. Furthermore, from the computation that established P was the midpoint of \overline{MN} we have $r_A r_B = PM \cdot AB = 17 \cdot 5 = 85$. As a result,

$$r_A + r_B = \sqrt{r_A^2 + r_B^2 + 2r_A r_B} = \sqrt{17^2 + 2 \cdot 17 \cdot 5} = \boxed{3\sqrt{51}}$$

7. In triangle ABC, points D, E, and F are on sides BC, CA, and AB respectively, such that BF = BD = CD = CE = 5 and AE - AF = 3. Let I be the incenter of ABC. The circumcircles of BFI and CEI intersect at $X \neq I$. Find the length of DX.

Proposed by Howard Halim

Answer: 3

Solution. The key observation is that X lies on BC with ID = IX. To prove the first claim, observe via angle chasing that


$$\angle IXC = \angle BFI = \angle BDI$$
,

where the last equality holds since BI is the perpendicular bisector of \overline{DF} . Similarly, $\angle IXB = \angle IDC$. Thus, $\angle BXI + \angle CXI = 180^{\circ}$, whence $X \in BC$. The claim IX = ID follows from the fact that I is the midpoint of, for example, arc \widehat{EX} . (The claim that X lies on BC also follows from Miquel's Theorem.)

Finally, remark that X and D are reflections across the tangency point T of the incircle with \overline{BC} , and so

$$DX = TC - TB = (s - c) - (s - b) = b - c$$

= $(AE + EC) - (AF + FB) = AE - AF = 3$.



8. Let \mathcal{E} be an ellipse with foci F_1 and F_2 . Parabola \mathcal{P} , having vertex F_1 and focus F_2 , intersects \mathcal{E} at two points X and Y. Suppose the tangents to \mathcal{E} at X and Y intersect on the directrix of \mathcal{P} . Compute the eccentricity of \mathcal{E} . (A parabola \mathcal{P} is the set of points which are equidistant from a point, called the focus of \mathcal{P} , and a line, called the directrix of \mathcal{P} . An ellipse \mathcal{E} is the set of points P such that the sum $PF_1 + PF_2$ is some constant P0, where P1 and P2 are the foci of P2. The eccentricity of P3 is defined to be the ratio P3.

Proposed by David Altizio

Answer: $\frac{2+\sqrt{13}}{9}$

Solution. Let T be the intersection point of the tangents to \mathcal{E} at X and Y; note that $T \in F_1F_2$ by symmetry and that by the problem statement T lies on the directrix ℓ of \mathcal{P} . Recall that by the reflection property of ellipses, TX is the external angle bisector of $\angle F_1XF_2$. Thus the (Exterior) Angle Bisector Theorem implies $\frac{F_2X}{F_1X} = \frac{F_2T}{F_1T} = 2$.

Denote by Q and R the projections of X onto ℓ and F_1F_2 respectively, and let $F_1X=x$ and $F_1F_2=d$; then $TR=QX=XF_2=2x$. The Pythagorean Theorem applied to triangles F_1XR and F_2XR implies

$$(2x)^{2} - x^{2} = F_{2}X^{2} - F_{1}X^{2} = F_{2}R^{2} - F_{1}R^{2} = (2d - 2x)^{2} - (2x - d)^{2}.$$

Simplifying yields $3x^2 = 3d^2 - 4xd$. Thus the ratio r := d/x satisfies the equality $3 = 3r^2 - 4r$, and so $r = \frac{2+\sqrt{13}}{3}$.


Finally, the eccentricity of \mathcal{E} is $\frac{d}{F_1X+F_2X}=\frac{d}{3x}=\boxed{\frac{2+\sqrt{13}}{9}}$

9. In triangle ABC, points M and N are on segments AB and AC respectively such that AM = MC and AN = NB. Let P be the point such that PB and PC are tangent to the circumcircle of ABC. Given that the perimeters of PMN and BCNM are 21 and 29 respectively, and that PB = 5, compute the length of BC.

Proposed by Howard Halim

Answer: $\frac{200}{21}$

Solution. Observe that $\angle BAC = \angle PBC$, so triangles $\triangle ANB$ and $\triangle CPB$ are similar to each other. It follows by spiral similarity that $\triangle BNP \sim \triangle BAC$. In turn, $\angle BPN = \angle BCN$, whence BNCP is cyclic. Analogous reasoning implies M lies on this circle as well.



Furthermore, triangles BNP and PCM are actually congruent, since they are both similar to $\triangle BAC$ and have common side BP = CP. This means BN = MP and CM = NP, so quadrilaterals BMNP and CNMP are both isosceles trapezoids. In turn, MN = BP = CP = 5 and quadrilateral AMPN is a parallelogram.

Since the perimeter of $\triangle AMN$ is 21, the perimeter of $\triangle ABC$ is $21 + 29 - 2 \cdot 5 = 40$. Hence triangles ANM and ABC are similar with a scale factor of $\frac{21}{40}$, implying

$$BC = MN \cdot \frac{40}{21} = 5 \cdot \frac{40}{21} = \boxed{\frac{200}{21}}.$$

10. Four copies of an acute scalene triangle \mathcal{T} , one of whose sides has length 3, are joined to form a tetrahedron with volume 4 and surface area 24. Compute the largest possible value for the circumradius of \mathcal{T} .

Proposed by Misha Ivkov and David Altizio

Answer: $\sqrt{4+\sqrt{3}}$


Solution. Denote the tetrahedron \mathcal{T} by ABCD, where we set BC = 3 without loss. The key idea is to unfold the tetrahedron into a net, transforming it into triangle PQR as shown below. This leads into a key lemma.

Lemma 1 (1999 AIME #15, etc.). The foot H of the perpendicular from D to plane ABC is the orthocenter of $\triangle PQR$.

Proof. Note that

$$BH^2 - CH^2 = BD^2 - DC^2 = BP^2 - PC^2$$
;

the first equality follows from the Pythagorean Theorem while the second comes from the equalities DB = BP and DC = CP. Thus the Perpendicularity Lemma tells us $PH \perp BC$, so H lies on the altitude from P. Applying this reasoning cyclically yields the desired.



Since the surface area of \mathcal{T} is 24, the area of $\triangle ABC$ is 6, and so via the formula $V=\frac{1}{3}bh$ we may deduce DH=2. Let X denote the intersection of PH with BC, i.e. the foot of the altitude from P to BC. By a similar area argument as above, we deduce PX=4, and via the definition of reflection, XD=4 as well. So $XH=\sqrt{4^2-2^2}=2\sqrt{3}$, implying $PH=4+2\sqrt{3}$.

Finally, let R denote the circumradius of $\triangle ABC$. Then the circumradius of $\triangle PQR$ is 2R. Thus

$$2(2R)^{2} = PH^{2} + RQ^{2} = (4 + 2\sqrt{3})^{2} + 6^{2} = 8(4 + \sqrt{3}),$$

so
$$R = \sqrt{4 + \sqrt{3}}$$

Note that this tetrahedron actually exists when \mathcal{T} has side lengths approximately 3, 4.021, and 4.763.

11. (Estimation) Gunmay picks 6 points uniformly at random in the unit square. If p is the probability that their convex hull is a hexagon, estimate p in the form 0.abcdef where a, b, c, d, e, f are decimal digits. (A convex combination of points x_1, x_2, \ldots, x_n is a point of the form $\alpha_1 x_1 + \alpha_2 x_2 + \cdots + \alpha_n x_n$ with $0 \le \alpha_i \le 1$ for all i and $\alpha_1 + \alpha_2 + \cdots + \alpha_n = 1$. The convex hull of a set of points X is the set of all possible convex combinations of all subsets of X.)

Proposed by Max Aires

Answer: 0.122327